

Contents

	Foreword	V
	List of Contributors	XIX
	Acknowledgments	XXI
1	Introduction	1
	<i>Peter Zwick</i>	
1.1	History of the Limnologische Flusstation Schlitz	1
2	The Breitenbach and Its Catchment	5
	<i>Jürgen Marxsen, Rüdiger Wagner, and Hans-Heinrich Schmidt</i>	
2.1	Study Area	5
2.2	The Stream and the Catchment	7
2.3	Sampling Sites	18
3	Environmental Characteristics	21
	<i>Hans-Heinrich Schmidt</i>	
3.1	Climate and Weather	21
3.1.1	Global Irradiance	21
3.1.1.1	Local Effects of Irradiance in the Catchment	24
3.1.2	Precipitation	24
3.1.3	Wind	27
3.1.4	Air Temperature	32
3.1.5	Discharge	34
3.1.5.1	Base Flow and Base Flow Index (BFI)	36
3.1.5.2	Falling Limb	38
3.1.5.3	Rising Limb	39
3.1.5.4	Flow Duration Curve	39
3.1.5.5	High Water	41
3.1.5.6	Bank-Full	43
3.1.5.7	Water Saturation of the Valley	44
3.1.5.8	Input Average	44
3.1.5.9	Low Water	44
3.2	Attempt to Define Substratum Dynamics	44

3.2.1	Cross-Section Measurements	46
3.2.2	Wet Stream Area	46
3.2.3	Water Temperature	47
3.2.3.1	Water Temperature in the Sediment	47
3.2.3.2	Water Temperature in the Stream	47
3.3	Chemistry	56
3.3.1	Buffering Capacity (ABC–Acid Binding Capacity)	57
3.3.2	pH	59
3.3.3	Oxygen	60
3.3.4	Soluble Inorganic Elements	64
3.3.4.1	Methods	64
3.3.4.2	Nitrogen	64
3.3.4.3	Phosphate	69
3.3.4.4	Silicate	70
3.3.4.5	Sulfate	71
3.3.4.6	Chloride	71
4	Detrital Energy Sources	73
4.1	Organic Matter in Streams	73
	<i>Jürgen Marxsen</i>	
4.2	Particulate Organic Matter	74
	<i>Jürgen Marxsen and Rüdiger Wagner</i>	
4.2.1	Particulate Organic Matter in Streams	74
4.2.2	POM Inputs to the Breitenbach	75
4.2.2.1	Measuring Periods and Methods	75
4.2.2.2	Seasonal Distribution of CPOM Inputs	76
4.2.2.3	Amounts of CPOM Inputs in Various Years and Stream Sections	78
4.2.2.4	Composition of CPOM Inputs	81
4.2.2.5	Total Inputs of POM	82
4.2.3	Standing Stocks of POM in the Breitenbach	83
4.2.4	Outputs of POM from the Breitenbach	83
4.3	Dissolved Organic Matter	84
	<i>Jürgen Marxsen</i>	
4.3.1	Dissolved Organic Matter in Streams	84
4.3.2	Concentrations and Dynamics of Dissolved Organic Matter in the Breitenbach	84
4.3.3	Inputs and Outputs of Dissolved Organic Matter to and from the Breitenbach	89
4.3.4	Composition of Dissolved Organic Matter in the Breitenbach: Carbohydrates	91
5	Primary Producers	99
	<i>Eileen J. Cox, Jürgen Marxsen, and Thomas G. Horvath</i>	
5.1	Primary Producers in Streams	99
5.2	The Communities	100

5.2.1	Algae	100
5.2.1.1	Algal Assemblages in Streams	100
5.2.1.2	Spatial and Temporal Patterns of Diatom Occurrences in the Breitenbach	101
5.2.1.3	Variation in Algal Biomass	108
5.2.1.4	Effects of Discharge on Benthic Diatom Assemblages	112
5.2.1.5	Microhabitats	115
5.2.1.6	Species Occurrences and Ecological Inferences	117
5.2.1.7	Impact of Grazers on Algal Assemblage and Biomass	122
5.2.1.8	General Conclusions from a Particular Ecosystem	122
5.2.2	Macrophytes	124
5.3	Primary Production	126
6	Bacteria and Fungi	131
	<i>Jürgen Marxsen</i>	
6.1	The Role of Bacteria and Fungi in Streams	131
6.2	The Communities	134
6.2.1	Bacterial Abundance and Biomass	134
6.2.1.1	Abundance in Different Habitats	134
6.2.1.2	Effects of Temperature and Discharge on Suspended Bacterial Abundance	139
6.2.1.3	Bacterial Biomass	141
6.2.2	Bacterial Community Composition	142
6.2.2.1	Methodological Approaches	142
6.2.2.2	Cultivation	144
6.2.2.3	Fluorescence <i>in situ</i> Hybridization	145
6.2.2.4	Temperature Gradient Gel Electrophoresis	151
6.2.2.5	Sequence Analysis	154
6.2.2.6	Water versus Sediment Bacterial Communities	161
6.2.2.7	Community Composition: Conclusions	162
6.2.3	The Fungal Community	162
6.3	Microbial Metabolic Activity	163
6.3.1	Approaches to Microbial Metabolic Activity Measurements in the Breitenbach	163
6.3.2	Extracellular Enzyme Activity	164
6.3.2.1	The Function of Extracellular Enzymes in Streams	164
6.3.2.2	Methodological Approach	166
6.3.2.3	Spatial Distribution	168
6.3.2.4	Activity of Different Enzymes	170
6.3.2.5	Environmental Controls and Regulation	175
6.3.2.6	The Molecular Biology Perspective	179
6.3.3	Bacterial Production	182
6.3.3.1	The Importance of Bacterial Carbon Production in Streams	182
6.3.3.2	Methodological Approach	183
6.3.3.3	Seasonal Fluctuation	184

6.3.3.4	Small-Scale Variability and Particulate Organic Matter Content	188
6.3.3.5	Influence of Temperature and Discharge	190
6.3.3.6	Comparing BCP between Different Streams and Rivers	191
6.3.3.7	Annual Bacterial Production and Respiration	193
6.3.4	Fungal Production	194
7	The Fauna of the Breitenbach	195
	<i>Peter Zwick, Georg Becker, Rüdiger Wagner, and Joachim Reidelbach</i>	
7.1	Sampling Methods	196
	<i>Peter Zwick</i>	
7.1.1	Collecting the Macrozoobenthos	196
7.1.1.1	Study of Specimens	197
7.1.1.2	Secondary Production	198
7.1.1.3	Life Cycle Analyses	201
7.1.1.4	Effect of an Insecticide Poisoning	201
7.1.2	Collecting Insects in Emergence Traps	202
7.1.2.1	Goals	202
7.1.2.2	Types of Trap Used and Years of Operation	202
7.1.2.3	Fundamentals of Emergence Trap Function	203
7.1.2.4	Errors during Trap Operation	205
7.1.2.5	Emergence Traps as Activity Traps and Terrestrial By-Catch	206
7.1.2.6	Emergence Trap Collections and Estimates of Secondary Production	207
7.1.2.7	Evaluation—Statistical Treatment of Data	208
	<i>Rüdiger Wagner</i>	
7.2	Protozoa: Ciliophora	209
	<i>Peter Zwick</i>	
7.3	Platyhelminthes and Annelida	211
	<i>Peter Zwick</i>	
7.3.1	Platyhelminthes	211
7.3.2	Annelida	212
7.4	Nematoda	212
	<i>Peter Zwick and Heino Christl</i>	
7.5	Mollusca	214
	<i>Peter Zwick</i>	
7.6	Arthropoda	216
	<i>Peter Zwick</i>	
7.6.1	Chelicerata	216
7.6.2	Hydrachnidia	217
7.6.3	Crustacea	218
7.6.4	Insecta	219
	<i>Peter Zwick</i>	
7.6.4.1	General and Minor Taxa	219
7.6.4.2	Ephemeroptera	226
7.6.4.3	Plecoptera	254

- 7.6.4.4 Coleoptera 309
- 7.6.4.5 Trichoptera 327
Rüdiger Wagner
- 7.6.4.6 Aquatic Diptera 369
Rüdiger Wagner
- 7.7 Chordata 404
Peter Zwick
- 7.8 Ecophysiology, Behavior, and Life Cycles of Scrapers 405
Georg Becker
- 7.8.1 Crowding Problems and Feeding Strategy of *Tinodes rostocki* (Trichoptera: Psychomyiidae), a Dominant Epilithic Caddisfly in the Middle and Lower Reaches of the Breitenbach 405
- 7.8.1.1 Larval Age Structure, Colonization of Natural Substrata, and Intraspecific Competition for Space in *T. rostocki* 405
- 7.8.1.2 Retreats of *T. rostocki* as a Substratum for Larval Food Resources 412
- 7.8.1.3 Feeding strategy of *T. rostocki* 419
- 7.8.2 Life Cycle and Ovipositing Adaptations of *Agapetus fuscipes* (Trichoptera, Glossosomatidae) to a First-Order Stream 421
- 7.8.2.1 Life Cycle of Glossosomatidae 421
- 7.8.2.2 Measurements of Pronotum Length, Larval Mass, and Case Length in *A. fuscipes* 422
- 7.8.2.3 Number of Larval Instars 423
- 7.8.2.4 Larval Mass and Case Length throughout the Ontogeny 424
- 7.8.2.5 Molt Increments 426
- 7.8.2.6 The Presence of Developmental Stages in the Breitenbach throughout the Year 427
- 7.8.2.7 Life Cycle Adaptation of *A. fuscipes* to a Small Uphill Stream 428
- 7.8.2.8 Ovipositing Behavior of *Agapetus fuscipes* 429
- 7.8.2.9 Egg Number and Egg Masses per Female 430
- 7.8.2.10 Distribution and Abundance of Egg Masses in the Breitenbach 430
- 7.8.2.11 Correlation between Egg Number and Cap Stone Size 431
- 7.8.2.12 Ovipositing Behavior of Glossosomatidae 432
- 7.8.3 Importance of Case Construction for Crawling Behavior and the Coexistence of Three Trichopteran Scrapers in the Upper Reach of the Breitenbach 432
- 7.8.3.1 Functions of Different Case Constructions for Trichopteran Species 432
- 7.8.3.2 Case Construction with Respect to Size, Shape, Composition, and Mass 433
- 7.8.3.3 Number and Size of Sand Grains 435
- 7.8.3.4 Use of Silk, Case Construction Costs, and Case: Larval Mass Quotient 436
- 7.8.3.5 Crawling Velocity of Larvae 436
- 7.8.3.6 Advantages of Different Case Constructions for Epilithic Scrapers 438

7.8.3.7	Effect of Substratum Roughness on Larval Crawling Velocity	439
7.8.4	Interactions between Scrapers and Periphyton	442
7.8.4.1	Introduction	442
7.8.4.2	Comparison of the Dietary Composition of Epilithic Trichopteran Species	445
7.8.4.3	Gut Content and Periphyton Analyses	446
7.8.4.4	Comparison between Species	446
7.8.4.5	Gut Content and Periphyton Composition throughout the Year	447
7.8.4.6	Variability in Gut Content and Periphyton Composition along the Stream	451
7.8.4.7	Primary Producers, Food Quality, and Resource Overlap	453
7.8.4.8	Interspecific Competition between Scrapers and Top-Down Control of Biofilms	455
7.8.4.9	Spatial and Temporal Development of Trichopteran and Ephemeropteran Grazers and Periphyton along the Breitenbach	459
7.8.4.10	Significance of Phototrophic Microorganisms for the Mortality, Growth, and Development of Trichopteran Scrapers	461
7.8.4.11	The Diurnal Foraging Behavior of <i>A. fuscipes</i> and <i>D. annulatus</i>	461
7.8.4.12	Food Choice in Laboratory Experiments	462
7.8.4.13	Response of Trichopteran Grazers to Biofilms of Diverse Quantity and Quality	467
7.8.4.14	Feeding Strategies and Ecological Segregation of Trichopteran Grazers in the Breitenbach	471
7.8.4.15	Conclusions	475
7.9	Effects of Water Temperature along the Breitenbach	476
	<i>Rüdiger Wagner</i>	
7.9.1	Water Temperature as an Important Variable for Explaining Body Size	476
7.9.2	Functional Feeding Groups	477
7.9.3	Experiments with <i>Chaetopteryx villosa</i>	478
7.9.3.1	Life Cycle and Adult Size at Different Stream Sites	478
7.9.3.2	Between-Sexes Size Difference	478
7.9.3.3	Field Experiment—Effects of Food and Temperature	480
7.9.3.4	<i>Chaetopteryx villosa</i> —Growth Models	481
7.9.4	<i>Gammarus fossarum</i> —Another Example	481
7.9.5	Water Temperature—A Limiting Resource	483
7.9.6	Scrapers—Indication of Site-Specific Intra- and Interspecific Competition Based on Emerged Adults	483
7.9.6.1	Caddisfly Scrapers—Size Differences along the Breitenbach	483
7.9.6.2	Within- and Between-Species Resource Limitation	484
8	Ecosystem Breitenbach	487
8.1	The Sandy Streambed Areas of the Breitenbach	487
	<i>Rüdiger Wagner, Jürgen Marxsen, and Hans-Heinrich Schmidt</i>	
8.1.1	Importance of Sediments in Streams	487

- 8.1.2 The Breitenbach Streambed 488
- 8.1.3 Interstitial Water: Chemical Characteristics 490
- 8.1.4 Breitenbach Sandy Sediments as Habitat 495
- 8.2 Aquatic Insects Emerging over an Artificially Changed Stream Bottom 496
Rüdiger Wagner
- 8.2.1 The “Silted” Emergence Trap 498
- 8.2.2 Laboratory Experiments 499
- 8.2.3 Emerging Aquatic Insects 499
- 8.3 Discharge Patterns Largely Determine Species Abundance and Community Diversity 503
Rüdiger Wagner and Hans-Heinrich Schmidt
- 8.3.1 Community and Variables Data Treatment 505
- 8.3.2 Environmental Variables 508
- 8.3.3 Community 512
- 8.3.4 Abundance and Discharge Pattern 514
- 8.3.5 Ordination 517
- 8.3.6 Discriminant Analysis 517
- 8.3.7 Community Diversity 519
- 8.3.8 Periods before and after 1990 522
- 8.3.9 The EPT Community of the Breitenbach and Environmental Variation 522
- 8.3.10 Multiple Stable Community States in Running Waters? 524
- 8.4 Eleven Years Emergence at Four Sites along the Breitenbach and the Effects of Discharge Patterns on the EPTD Community 526
Rüdiger Wagner and Hans-Heinrich Schmidt
- 8.4.1 Variables, Community, and Statistics 527
- 8.4.2 Environmental Variables 528
- 8.4.2.1 Precipitation 528
- 8.4.2.2 Discharge 528
- 8.4.2.3 Water Temperature 529
- 8.4.2.4 Air Temperature and Number of Rainy Days 529
- 8.4.3 Community 529
- 8.4.3.1 Distribution of Species along the Breitenbach 529
- 8.4.3.2 Species Affected by Discharge Pattern 531
- 8.4.3.3 Species with Simultaneous Spatial and Temporal Variability 534
- 8.4.3.4 Correspondence Analysis 534
- 8.4.3.5 Community Measures 537
- 8.4.3.6 Discriminant Analysis 539
- 8.4.4 Discussion 539
- 8.5 The Erlenbach—a Comparison with the Breitenbach 543
Rüdiger Wagner
- 8.5.1 Environmental Variables 545
- 8.5.1.1 Conductivity and pH 545
- 8.5.1.2 Discharge 545

8.5.1.3	Air and Water Temperature	545
8.5.2	Community	547
8.5.3	Conclusion	553
8.6	Modeling Discharge and Insect Abundance with Artificial Neural Networks (ANNs)	554
	<i>Michael Obach and Rüdiger Wagner</i>	
8.6.1	Precipitation–Discharge	555
8.6.2	Prognosis of Yearly Abundance of 17 EPT Species	557
8.6.3	Seasonal Effects of Variables on Species in Models	559
8.6.4	Effects of Discharge Parameters on Predictions of Monthly Abundances of <i>Baetis rhodani</i>	560
8.6.5	Predictions of Yearly Abundance Patterns	561
8.6.6	Abundance Predictions for the Caddisfly <i>Apatania fimbriata</i> at Four Sites along the Stream	562
8.6.7	General Aspects	563
8.7	Food Chains and Carbon Flow	564
	<i>Jürgen Marxsen</i>	
8.7.1	The Organic Matter Budget of the Breitenbach	564
8.7.2	Bacteria in the Food Web System of the Breitenbach	567
8.8	Global Change and Timing of Insect Emergence at the Breitenbach	570
	<i>Peter Zwick and Rüdiger Wagner</i>	
8.8.1	Introduction	570
8.8.2	Methods	572
8.8.2.1	Selection Criteria	572
8.8.2.2	Accuracy of Emergence Data and Possible Sources of Error	573
8.8.2.3	Shift in Emergence Dates	573
8.8.2.4	Emergence Dates and Temperature Regimes	573
8.8.3	Results	575
8.8.3.1	Shift in Emergence Time of Individual Ephemeroptera and Plecoptera Species	575
8.8.3.2	Shift in Emergence Time of Trichoptera Species	576
8.8.3.3	Emergence Dates and Thermal Regime	577
8.8.4	Discussion	578
9	Summary	585
	<i>Georg Becker, Eileen J. Cox, Jürgen Marxsen, Hans-Heinrich Schmidt, Rüdiger Wagner, and Peter Zwick</i>	
9.1	Climate, Environment	585
9.1.1	Hydrological Chain	586
9.1.2	Thermal Chain	586
9.1.3	Water Chemistry	587
9.2	Detrital Energy Sources	587
9.3	Primary Producers	588
9.4	Bacteria	589

9.5	Organic Matter Balance and Carbon Flow	590
9.6	Fauna	591
9.6.1	The Dominant Pterygota	591
9.6.1.1	Ephemeroptera	592
9.6.1.2	Plecoptera	593
9.6.1.3	Trichoptera	594
9.6.1.4	Diptera	595
9.6.1.5	Coleoptera	595
9.6.2	Scrapers of the Breitenbach Community	596
9.7	Ecosystems	598
9.7.1	The Stream Bottom	598
9.7.2	Ordination	598
9.7.3	Modeling	599
9.7.4	Erlenbach	599
9.7.5	Global Change	600
	References	601
	Index	663

