

[illegible]

This is a title figure. The positions of the figure and the legend depend on the amount of text and of the size of the figure. The final position will be fixed by the typesetters.

© 2010 by Astro Ltd.

Published exclusively by WILEY-VCH Verlag GmbH & Co. KGaA

Demonstration of the overall layout of the *Laser Physics Letters* L^AT_EX class

*First Author*¹ and *Second Author*^{2,*}

¹ First Institute² Second Institute

Received: XXXX, Accepted: XXXX

Published online: XXXX

Key words: List; of; semicolon; separated; keywords

1. First section of your article

The dramatic advances of quantum- and atom-optics allow at present accurate control over the internal and external degrees of freedoms of atoms. In particular, atoms can be routinely cooled down to temperatures in the nK range, they can be stored in magnetic or optical traps for up to several minutes and a precise control over their quantum state is possible. These achievements motivate several new directions, such as the implementation of quantum information processing using neutral atoms or the construction of matter-wave interferometers.

Besides these final goals, experiments investigating the coherent quantum behaviour of neutral atoms provide new insights into quantum physics and into new applications of quantum technology in a rapidly expanding area of science and technology. Engineering quantum systems with

neutral atoms require means for coherent manipulation and state sensitive detection.

A natural way to attempt coherent control over atoms is using potential structures which are comparable in size with the de Broglie wavelength of the atom. Optical standing wave fields, so called optical lattices, generate potential structures on this length scale, typically on the order of the optical wavelength ($\sim \lambda/2$), and are sufficiently strong to confine and structure an ultracold cloud of atoms. Alternatively, optical micro traps using diffractive and refractive elements and magnetic micro traps, based on micro-fabricated magnets have been introduced. In micro traps, the potential is defined by the geometry of the microstructure and almost arbitrary spatial and temporal potential shaping is possible. The implementation of optical, electrostatic, electrodynamic and magnetic micro potentials on a single chip not only promises a versatile control over

* Corresponding author: e-mail-Address




Figure 1 A figure over one column

atomic quantum states but also makes widespread applications possible. Matter wave interferometers using micro traps have recently been realized [2,4]. Interference fringes related to the quantum mechanical phase of atomic Bose-Einstein condensates were thereby detected by standard absorption imaging. This is possible due to the large number of atoms involved in the interference. However, quantum information processing or interferometry with single atoms require the development of detectors, sensitive to the signal of single atoms. At present, research is focussed on both coherent manipulation techniques and state selective single atom detection. Some aspects of the recent progress on coherent manipulation using micro traps based on micro-fabricated current-carrying and optical structures are reviewed in this article. The selection is far from being complete. The tremendous growth of the field during the last years does not allow us to present a comprehensive overview and we are forced to restrict ourselves to describe an admittedly biased selection.

2. Second section

Quantum information processing with neutral atoms offers a set of advantages over other approaches:

(1) Flexible implementation of qubits: atoms and molecules as carriers of quantum information represent intrinsically identical systems of qubits which can be decoupled from their environment to a high degree. In addition, there exist a range of possibilities to encode quantum information in quasi spin-1/2 systems. For atoms in micro-traps quantum information can be encoded in the internal (e.g. Hyperfine ground states) as well as external degrees of freedom (e.g. vibrational modes of the trapping potential).

(2) Scalability: trapping geometries based on magnetic and optical micro-traps (Fig. 1) allow for the implementation of hundreds or even thousands of identical atomic qubits in parallel. Based on optical lattices the number of qubits

can be increased to about 10^5 . This gives the unique possibility to develop and experimentally test schemes for quantum information processing on large scales.

3. Further section of your article

This is the text of this article. More text, much more text,
and even more text. This is the text of this article. More
text, much more text, and even more text. This is the text of
this article. More text, much more text, and even more text.
This is the text of this article. More text, much more text,
and even more text. This is the text of this article. More
text, much more text, and even more text. This is the text
of this article. More text, much more text, and even more
text. This is the text of this article. More text, much more
text, and even more text. This is the text of this article.
More text, much more text, and even more text. This is the
text of this article. More text, much more text, and even
more text. This is the text of this article. More text, much
more text, and even more text.

This is the text of this article. More text, much more text, and even more text. This is the text of this article. More text, much more text, and even more text. This is the text of this article. More text, much more text, and even more text. This is the text of this article. More text, much more text, and even more text. This is the text of this article. More text, much more text, and even more text.

Some lists:

- First entry
 - use the `{itemize}` environment
 - Second entry
-
1. First entry with a number
 - use the `{enumerate}` environment
 2. Second entry with a number
 - itemize within enumerate
 - First entry without number within a list with numbers
 - Second entry without number within a list with numbers

3.1. Change marks

Please use for changes requested by the referee the following colour change option: **This is a text snippet marked as *changed*.** This is done by enclosing it in an environment called `changed`. Please note that in certain circumstances there might be small side effects such as make up deviations or additional blanks.

Acknowledgements Insert your acknowledgements here.




Figure 2 A figure wider than 1 column.
Use the `\beginfigure*` command.

Making bibliographies using BibTeX and lpl.bst

At the end of this file you see an example bibliography which represents the journals reference style [1–6]. Please format your real bibliography accordingly.

Optionally, you may generate your bibliography using BibTeX, with the bib-style-file `pss.bst` from the template package in your L^AT_EX search path. To this end, replace the example database-file `lpl-demo.bib` with your own existing one, or alternatively use it as a template for generating your new database. The following code in your manuscript source file enables BibTeX-functionality:

```
\bibliographystyle{lpl}
\bibliography{<database-filename>}
```

You then need to run your manuscript source file through BibTeX using the command

```
bibtex <filename>
```

(most T_EX frontends have a shortcut for this) and afterwards through L^AT_EX twice, in order to get the correct label numbering.

Important: Before sending your manuscript source file to the publisher, remember to transfer the actual, fully formatted bibliography contained in the BibTeX-generated file `<filename>.bb1` to your manuscript source file.

References

- [1] F. M. Firstauthorfamilyname, F. M. Secondauthorfamilyname, and C. Lastauthorfamilyname, *Abbreviatedjournal-name* **volume**, page (year).
- [2] F. Exemplename and I. E. Anotherauthorname, *phys. stat. sol. (a)* **1**, 111 (2050).
- [3] A. Firstauthorname, B. Secondauthorname, and C. Thirdauthorname, *Here Goes the Title of the Book* (Publisher, City, year), p. 111.
- [4] A. Firsteditorname, B. Secondeditorname, and C. Thirdeditorname (eds.), *Here Goes the Title of the Edited Book* (Wiley-VCH, Berlin, 2050), p. 111.
- [5] D. Contributorname, in: *The Title of the Book*, edited by The Name of the Editors, Followed by the Title of the Series of Books (Publisher, City, year), chap. 1.
- [6] A. Lastbutnotleastname, *Proceedings 1st Dummy Conference on Citation Formatting*, City, Country, Part A (Publisher, City, year), pp. 1–11.